

Citation: Yenice MG, Seker KG, Sam E, et al. Comparison of cold-knife optical internal urethrotomy and holmium:YAG laser internal urethrotomy in bulbar urethral strictures. *Cent European J Urol.* 2017; doi: 10.5173/ceju.2017.1391 [Epub ahead of print]

Comparison of cold-knife optical internal urethrotomy and holmium:YAG laser internal urethrotomy in bulbar urethral strictures

[Autor's unedited version]

Mustafa Gurkan Yenice, Kamil Gokhan Seker, Emre Sam, Yunus Colakoglu, Feyzi Arda Atar, Selcuk Sahin, Abdulmuttalip Simsek, Volkan Tugcu

Department of Urology, Bakirkoy Dr.Sadi Konuk Training and Research Hospital, Turkey, Istanbul

Key Words: urethral stricture, holmium laser, male urethra

Corresponding author

Kamil Gokhan Seker
gkhseker@hotmail.com

Article history

Submitted: April 20, 2017

Accepted: Nov. 13, 2017

Published online: Dec. 6, 2017

Introduction To compare the results of cold-knife optical internal urethrotomy (OIU) and Holmium:YAG laser internal urethrotomy (HIU) in primary bulbar urethral strictures.

Material and methods A total of 63 patients diagnosed with primary bulbar urethral stricture between August 2014 and September 2015 were assigned to the OIU (n = 29) and HIU (n = 34) groups. The demographic variables, biochemistry panels, and preoperative and postoperative uroflowmetry (UF) results including the maximum flow rate (Q_{max}) and mean flow rate (Q_{mean}) values, retrograde urethrography, and diagnostic flexible urethroscopy findings were recorded prospectively. Demographic features and preoperative values were not statistically different between groups (p >0.05). Mean surgical times were 18.4 +/-2.3 min for OIU and 21.9 +/- 3.8 min for HIU groups, which was statistically significant (p <0.05). There was no significant difference in complication rates in both groups (p = 0.618).

Results Postoperative Q_{max} values were increased in both groups even though postoperative Q_{max} values were not significantly different between the two groups in the short- and long-term results at 3, 6, and 12 months (p >0.05). There was no recurrence in the first 3 months in both groups. The urethral stricture recurrence rate up to month 12 was not statistically significant for the OIU group (n = 6, 20.7%) compared with the HIU group (n = 11, 32.4%; p = 0.299).

Conclusions Holmium:YAG laser internal urethrotomy is an alternative method to cold-knife optical internal urethrotomy, and it has similar success rates in the treatment of short segment bulbar urethral strictures.

INTRODUCTION

Urethral strictures are still among the most common reasons of urology clinic admissions. In addition to their negative impact on quality of life, urethral strictures are also associated with further financial burden related to urinary tract infections, bladder stones, fistulas, sepsis and possibly renal insufficiency [1].

Minimally invasive surgical approaches are utilized efficiently and safely in the treatment of various disorders in urology practice, treatment options for urethral strictures have been limited both in number and level of success, as judged by the high recurrence rates. Treatment options for urethral strictures include self-catheterization, bougie dilation, balloon dilation, urethral stent placement, cold-knife optical internal urethrotomy (OIU), laser urethrotomy methods and open reconstructive urethroplasty, which is generally reserved for cases who did not benefit from endoscopic approaches [2]. OIU, described by Sachse in 1974, is still the preferred method due to the practical and minimally invasive nature of the procedure [3]. One of the alternative approaches is the correction of the stricture with laser energy, which was first used in urology in 1984 and established as a common method in internal urethrotomy since the nineties [4]. In the past, several prospective randomized clinical trials comparing cold incision internal urethrotomy with holmium yttrium-aluminium-garnet (Ho:YAG) laser urethrotomy have been reported [4-9].

The aim of this prospective randomized study is to compare pre- and post-operative uroflowmetry results and operations characteristics of OIU and Ho:YAG laser approaches in bulbar urethral strictures.

MATERIAL AND METHODS

All patients diagnosed with primary bulbar stricture and underwent internal urethrotomy (IU) in our clinic between August 2014 and September 2015 were enrolled in the study. Local review board approved study protocol. Patients with recurrent urethral strictures, a history of internal urethrotomy, abnormal ultrasound findings in the bladder (e.g. bladder tumor / calculi), suprapubic cystostomy, non-bulbar strictures and those with a stricture segment longer than 2 cm were excluded from the study. Urethral stricture was diagnosed with history, uroflowmetry, ultrasound, retrograde urethrography and diagnostic flexible urethroscopy performed under local anesthesia. Complete blood count, biochemistry panel and urinalysis was evaluated in all patients. Coagulation times were corrected in the patients with abnormal coagulation test results and urinary tract infections were duly treated with antibiotics and sterile urine was attained prior to the operation. The patients were randomly assigned to one of the two treatment groups, cold-knife OIU (group 1) and Ho:YAG laser internal urethrotomy (HIU) (group 2). All patients were evaluated with ultrasound, uroflowmetry, international prostate symptom score (IPSS), quality of life (QoL) results preoperatively and at 3rd, 6th and 12th postoperative months. The procedure was accepted as successful when the patient did not complain of any persistent subjective and objective symptoms and the Qmax was >12 ml/s. Diagnostic flexible urethroscopy was performed in patients with persistent subjective and objective voiding symptoms or/and that Qmax was lower than 12 ml/sn. If urethral strictures was detected by flexible cystoscopy, these was accepted as recurrent strictures.

All operations were performed in the lithotomy position. Prophylactic intravenous cefuroxime axetil 750 mg was administered with the induction of anesthesia. The guidewire (Boston Scientific, USA) was advanced to the urethra through the stricture with the help of a 7.5 French Wolf ureterorenoscope (Richard Wolf, Knittlingen, Germany) in all patients before the intervention (Figure 1, 2). Cold-knife OIU was performed with a straight type urethrotomy knife inserted through a 21 Charrière Storz

Citation: Yenice MG, Seker KG, Sam E, et al. Comparison of cold-knife optical internal urethrotomy and holmium:YAG laser internal urethrotomy in bulbar urethral strictures. *Cent European J Urol.* 2017; doi: 10.5173/ceju.2017.1391 [Epub ahead of print]

urethrotome (Karl Storz, Germany). Laser urethrotomy was performed with a Ho: YAG laser litho fiber (LISA/Sphinx 30) with the frequency set between 4 – 20 Hz, power set at 30 W and pulse energy of 0.5–4.0 J. Both interventions were aimed at 12 o'clock position (Figure 3, 4). A 18 F foley urethral catheter was placed over the guidewire postoperatively and removed 3 days later. Patients with suspicion of stricture recurrence was performed 16 French flexible cystoscopy (Karl Storz, Germany).

A total of 63 patients aged between 26 and 69 years (mean age +/- standard deviation: 55.1 +/-9.1 years) were included in the study. There was no significant difference between the mean ages of the treatment groups, which were 54.8 +/-9.5 years for OIU (n = 29) and 55.3 +/-8.9 years for HIU (n = 34) (p = 0.895). Of the 63 patients, 59 (93.7%) were married, with 4 single patients in each treatment group. Urethral stricture was iatrogenic in 42 (66.7 %) patients, secondary to trauma in 12 (19%), secondary to urethritis in 6 (9.5%) and idiomatics 3 (4.8%) patients. There were no significant differences between the mean age, marital status and preoperative Qmax values between the two groups (p <0.05 for all comparisons). The demographic features and etiology of the strictures are present in Table 1.

SPSS 22.0 program (SPSS Inc; Chicago, IL, USA) was used in the analyzes. The distribution of the variables was measured by the Kolmogorov simirnov test. Mann-Whitney U test was used in the analysis of quantitative independent data. Chi-square test were used for the analysis of independent data and Fischer test was used when Chi-square test conditions were not met. Statistical significance was set at a p value of <0.05.

RESULTS

OIU group and HIU group did not difference significantly (p >0.05) in Q max values for preop, postop 3 months, postop 6 months, postop 12 months. Flexible urethroscopy was performed 25 patients that with objective voiding symptoms and/or Qmax <12 m/s. Flexible urethroscopy detected recurrence in 6 (20.7%) patients in OIU and 11 (32.4%) patients in HIU (p = 0.299) who were symptomatic and had declining Qmax values at 12th postoperative month (Table 2).

The mean operative time in HIU, 21.9 +/-3.8 min, was significantly longer compared to OIU (mean operative time 18.4 +/-2.3 min) (p <0.05) (Figure 5) .Three patients in OIU groups underwent bleeding after urethrotomy and one patient in HIU groups underwent urine extravasation postoperatively. All complications were managed conservatively.

Preoperative overall mean Qmax was 6.5 +/-0.7 mL/sec, with no significant difference between the treatment groups (6.5 +/-0.7 mL/sec for OIU and 6.5 +/-0.7 mL/sec for HIU) (p = 0.874).

DISCUSSION

Although blunt perineal trauma, urethral instrumentation, lichen sclerosis of atrophicus, sexually transmitted diseases and a history of urethral infections are listed among the underlying reasons for urethral strictures, the most cases observed in clinical practice are iatrogenic [10]. First detectable histological changes are squamous metaplasia of the pseudostratified epithelium into columnar cells, resulting in loss of fluid isolation and increase in permeability which leads to urine extravasation and fibrosis [11]. Major factors to be considered in treatment selection are etiology, stricture length, depth of scar and localization [12, 13]. The etiology of the patients in our study group was consistent with the previous reports with majority of patients classified as iatrogenic (n= 42, 66.7 %), while 12 patients (19%) had a stricture secondary to trauma, 6 patients (9.5 %) related to a history of urethritis and 3 patients (4.8 %) had a idiomatic. In addition, all strictures were localization in bulbar urethra and having short-segment bulbar urethral stricture (less than 2 cm).

Citation: Yenice MG, Seker KG, Sam E, et al. Comparison of cold-knife optical internal urethrotomy and holmium:YAG laser internal urethrotomy in bulbar urethral strictures. *Cent European J Urol.* 2017; doi: 10.5173/ceju.2017.1391 [Epub ahead of print]

Various alternative approaches have been proposed for urethral stricture treatment over the years and there is no consensus on the most successful method yet. Along with advances in laser technology, laser urethrotomy is widely used. Most common laser types used for urethral stricture treatment are neodymium-doped yttrium aluminium garnet (Nd: YAG), argon, potassium titanyl phosphate (KTP), thulium and Ho: YAG [14, 15]. In this study, we used; a Ho: YAG laser with the frequency set between 4–20 Hz, power set at 30 W and pulse energy of 0.5–4.0 J.

OIU is a more practical and economical approach compared to HIU whereas high recurrence rates and bleeding rates. [6]. One of the advantages of laser urethrotomy over OIU is the scar tissue that remains after the incision which is very useful in haemostasis. Laser approach also provides a better visualization of the lesion but unfortunately this method is not widely available due to relatively higher costs [16, 17].

Previous studies have reported contradicting results regarding the duration of the procedure for OIU and HIU. While Atak et al [5] reported a significantly shorter mean operative time for HIU (16.4 +/- 8.04 min) compared to OIU (23.8 +/-5.47 min), performing the OIU (7.44 min) took significantly shorter HIU (19.8 min) in the report by Jain et al [8]. In our study, similar to Jain et al. study, the duration of operation was statistically longer in HIU group according to the OIU group. The difference between these results may be related to technical difficulty and lack of experience for laser treatment.

The most widely accepted method for minimizing bleeding in internal urethrotomy is incising the scar tissue at 12 o'clock position between the corpus cavernosum or wherever the scar tissue is more extensive. Kamp et al. compared the Ho:YAG laser incisions at 12 and 6 o'clock positions and could not detect a difference between the complication or bleeding rates between the groups [7]. In a series of 190 patients treated with HIU, surgery related complications were seen in 23 patients but it was noted that nearly half of the complications occurred in the first 50 procedures and all were resolved with conservative measures [6]. In another study, Jhanwar et al. reported a total complication rate of 14.6%. In addition to our study they reported fever and urinary tract infection [9]. We performed all the incisions at the 12 o'clock position and did not have major complications or cavernous body injury. Bleeding was detected in 3 patients in the OIU group and treated by perineal compression. Bleeding was not encountered in HIU group. This condition may be explained by the fact that the Ho: YAG laser can provide coagulation.

In a randomized clinical study reported by Atak et al, success rates at 12 months for HIU and OIU were 81% and 53%, respectively [5]. A meta-analysis of 44 articles comprising 3230 patients reported the success rates in urethral strictures as 74.9% for laser and 68.5% for cold- knife urethrotomy [18]. In our study, The mean success rates for groups OIU and HIU were 79.3 % and 67.6%, respectively. We did not detect any differences between the success rates in both groups at all follow ups. Our results can be accepted like a similar study which compared laser versus cold- knife urethrotomy under local anesthesia [9].

Repetitive endoscopic interventions in bulbar urethral strictures are related to increases in stricture segment length and complexity and delay in urethroplasty operations [19]. Endoscopic treatment of urethral strictures are associated with high recurrence rates, ranging between 30% and 80% [20], which led to numerous studies comparing the efficacy of various treatment approaches . Sachse et al reported a recurrence rate of 23% in their first case series [21]. Comparable recurrence rates are still reported up to this day in urethral stricture treatment. Pansadoro and Emiliozzi reported a recurrence rate of 58% following OIU in bulbar urethral strictures [22]. Recurrences usually occur within 3-12 months, as Naude and Heyns emphasize [23]. In another study, Heyns et al. reported that recurrences within the first 3 months were a poor prognostic factor [24]. In our study, there was no recurrence in

Citation: Yenice MG, Seker KG, Sam E, et al. Comparison of cold-knife optical internal urethrotomy and holmium:YAG laser internal urethrotomy in bulbar urethral strictures. *Cent European J Urol.* 2017; doi: 10.5173/ceju.2017.1391 [Epub ahead of print]

the third month in both groups. When recurrence rates at 6 and 12 months were considered, recurrence was detected in 18 patients in both groups.

Another type of energy for urethrotomy is plasmakinetic. There are several studies in the literature about the treatment of urethral strictures with plasma kinetics [25, 26]. Until now, there is no clear criterion to assess success or failure after urethrotomy. Cecen at al. [25] and Ozcan at al. (26) were accepted to successful at their study, when the patient did not complain of any voiding difficulty and the Qmax was >12 ml/s. Another studies were accepted as successful if the patient did not report any voiding difficulty, the maximum flow rate (Qmax) was greater than 10 mL/sec [5, 9]. Tom at al. have reported that both sensitivity of Qmax and IPSS to detecting stricture recurrence is low. There is a need for tools with high sensitivity and specificity for urethral stricture recurrence follow-up [27, 28]. In this study, the procedure was accepted as successful when the patient did not complain of any persistent subjective and objective symptoms and the Qmax was >12 ml/s.

This study has several limitations. First, only short segment bulbar urethral strictures can be considered. Another limitations are that operations were performed by different surgeons and the lack of the questionnaire for assessment of urinary symptoms. High recurrence rates of urethral strictures following internal urethrotomy have incited the proposal of novel approaches for the treatment of this condition. Results with HIU have been encouraging so far but more improved approaches either with Ho:YAG or other novel methods will be necessary to achieve higher success and lower recurrence rates.

CONCLUSIONS

Urethral strictures are associated with a high burden on the quality of life and healthcare expenditures similar to other chronic disorders. Although various different methods have been proposed, none of them were able to prevent recurrences definitively. Lack of a significant difference between recurrence rates and complication rates of HIU and OIU suggests that both are safe and effective treatment methods in bulbar urethral strictures. Larger studies encompassing cases with different characteristics may help to identify the variables that may favor or discourage the use of either method in future studies.

Conflicts of interest

The authors declare no conflicts of interest.

Informed consent was obtained from all individual participants included in the study.

Citation: Yenice MG, Seker KG, Sam E, et al. Comparison of cold-knife optical internal urethrotomy and holmium:YAG laser internal urethrotomy in bulbar urethral strictures. *Cent European J Urol.* 2017; doi: 10.5173/ceju.2017.1391 [Epub ahead of print]




Figure 1. Endoscopic appearance of short segment bulbar urethral stricture.



Figure 2. Appearance of guidewire and urethral stricture segment.

Citation: Yenice MG, Seker KG, Sam E, et al. Comparison of cold-knife optical internal urethrotomy and holmium:YAG laser internal urethrotomy in bulbar urethral strictures. *Cent European J Urol.* 2017; doi: 10.5173/ceju.2017.1391 [Epub ahead of print]



Figure 3. Appearance of laser urethrotomy with a Ho: YAG laser litho fiber at 12 o'clock position.




Figure 4. Postoperative appearance of urethral lumen.

Citation: Yenice MG, Seker KG, Sam E, et al. Comparison of cold-knife optical internal urethrotomy and holmium:YAG laser internal urethrotomy in bulbar urethral strictures. *Cent European J Urol.* 2017; doi: 10.5173/ceju.2017.1391 [Epub ahead of print]




Figure 5. Operation time of OIU and HIU in both groups ($p < 0.05$ and $p < 0.05$, respectively).

Table 1. Patient's characteristics

	Min	Max	Median	Mean	± s.d./n-%
Age	26	69	57	55.1	± 9.1
Marital Status	Married			4	6.3%
	Single			59	93.7%
Etiology	Iatrogenic			42	66.7%
	Traumatic			12	19.0%
	Urethritis			6	9.5%
	Idiopathic			3	4.8%
Operation Time	15	30	19	20.3	± 3.6
Complication				4	6.3%
Qmax ml/sn					
Preoperative	5.1	7.9	6.4	6.5	± 0.7
3rd month	14.5	19.1	17.3	17.0	± 1.4
6th month	10.7	17.9	15.7	15.0	± 1.9
12nd month	8.9	15.9	13.9	13.3	± 2.1
Recurrence					

Citation: Yenice MG, Seker KG, Sam E, et al. Comparison of cold-knife optical internal urethrotomy and holmium:YAG laser internal urethrotomy in bulbar urethral strictures. *Cent European J Urol.* 2017; doi: 10.5173/ceju.2017.1391 [Epub ahead of print]

3rd month	0	0.0%
6th month	9	14.3%
12nd month	17	27.0%

Table 2 Comparison of the preoperative and postoperative parameters.

		Group I		Group II		P	
		Mean \pm s.d./n-%	Median	Mean \pm s.d./n-%	Median		
Age		54.8 \pm 9.5	58.0	55.3 \pm 8.9	57.0	0.895	m
MaritalStatus	Married	2	6.9%	2	5.9%	0.869	x ²
	Single	27	93.1%	32	94.1%		
Etiology	Iatrogenic	21	72.4%	21	61.8%		
	Traumatic	5	17.2%	7	20.6%		
	Urethritis	2	6.9%	4	11.8%		
	Idiopathic	1	3.4%	2	5.9%		
Operation Time		18.4 \pm 2.3	18.0	21.9 \pm 3.8	20.0	0.000	m
Complication		1	3.4%	3	8.8%	0.618	x ²
Qmax ml/sn							
Preoperative		6.5 \pm 0.7	6.4	6.5 \pm 0.7	6.4	0.874	m
3rd month		17.0 \pm 1.4	17.2	16.9 \pm 1.4	17.4	0.719	m

Citation: Yenice MG, Seker KG, Sam E, et al. Comparison of cold-knife optical internal urethrotomy and holmium:YAG laser internal urethrotomy in bulbar urethral strictures. *Cent European J Urol.* 2017; doi: 10.5173/ceju.2017.1391 [Epub ahead of print]

6th month	15.3	± 1.7	15.4	14.7	± 2.0	15.7	0.232	m
12nd month	13.7	± 2.0	14.1	13.0	± 2.1	13.8	0.159	m
Recurrence								
3rd month	0	0.0%		0	0.0%	-		X ²
6th month	3	10.3%		6	17.6%		0.409	X ²
12nd month	6	20.7%		11	32.4%		0.299	X ²

^mMann-whitney u test / ^{X²} Chi-square test (Fischer test)

Reference

1. Santucci RA, Joyce GF, Wise M. Male urethral stricture disease. *J Urol.* 2007; 177: 1667-1674.
2. Heyns CF, van der Merwe J, Basson J, van der Merwe A. Treatment of male urethral strictures possible reasons for the use of repeated dilatation or internal urethrotomy rather than urethroplasty. *S Afr J Surg.* 2012; 50: 82- 87.
3. Sachse H. Treatment of urethral stricture: transurethral slit in view using sharp section. *Fortschr Page 5/11 Med.* 1974;92:12-15.
4. Dutkiewicz SA, Wroblewski M. Comparison of treatment results between holmium laser endourethrotomy and optical internal urethrotomy for urethral stricture. *Int Urol Nephrol.* 2012; 44:717-724.
5. Atak M, Tokgoz H, Akduman B, Erol B, Donmez I, Hanci V, et al. Low-power holmium:YAG laser urethrotomy for urethral stricture disease: comparison of outcomes with the cold-knife technique. *Kaohsiung J Med Sci.* 2011;27:503-507.
6. Liu Q, Ma W, Li X, Zhang W, Cao W, Zhou Q, et al. Holmium Laser Endourethrotomy in Urethral Strictures. *Urol J.* 2014;11:1264-1270.
7. Kamp S, Knoll T, Osman MM, Köhrmann KU, Michel MS, Alken P. Low-power holmium: YAG laser urethrotomy for treatment of urethral strictures: functional outcome and quality of life. *J Endourol.* 2006; 20:38-41.
8. Jain SK, Kaza RM, Singh BK. Evaluation of holmium laser versus cold knife in optical internal urethrotomy for the management of short segment urethral stricture. *Urol Ann.* 2014; 6:328-333.
9. Jhanwar A, Kumar M, Sankhwar SN, Prakash G. Holmium laser vs. conventional (cold knife) direct visual internal urethrotomy for short-segment bulbar urethral stricture: Outcome analysis. *Can Urol Assoc J.* 2016; 10(5-6): E161-E164
10. Tritschler S, Roosen A, Füllhase C, Stief CG, Rübber H. Urethral stricture: etiology, investigation and treatments. *Dtsch Arztebl Int.* 2013;110:220-226.
11. Andrich DE, Mundy AR. Urethral strictures and their surgical treatment. *BJU Int.* 2000;86:571-580.
12. Singh O, Gupta SS, Arvind NK. Anterior urethral strictures:A brief review of the current surgical treatment. *Urol Int.* 2011; 86:1-10.
13. Wein AJ, Kavoussi LR, Novick AC, Partin AW, Peters CA. *Campbell- Walsh Urology.* 10th ed. Philadelphia, Elsevier Saunders, 2012, pp. 967-973
14. Hampson LA, McAninch JW, Breyer BN. Male urethral strictures and their management. *Nat Rev Urol.* 2014;11:43-50.
15. Wang L, Wang Z, Yang B, Yang Q, Sun Y. Thulium laser urethrotomy for urethral stricture: a preliminary report. *Lasers Surg Med.* 2010; 42: 620-623.
16. Dhabalia JV, Nelivigi GG, Punia MS, Kumar V. Refractory post visual internal urethrotomy bleeding managed by angioembolization. *Indian J Radiol Imaging.* 2010; 20:37-38.
17. Moreno Sierra J, Prieto Nogal S, Galante Romo I, Fernández Montarroso L, López Corral JC, Silmi Moyano A. The usefulness of laser in the treatment of urethral stenosis. *Arch Esp Urol.* 2008;61:978-984.
18. Jin T, Li H, Jiang LH, Wang L, Wang KJ. Safety and efficacy of laser and cold knife urethrotomy for urethral stricture. *Chin Med J (Engl)* 2010; 123:1589-1595.
19. Hudak SJ, Atkinson TH, Morey AF (2012) Repeat transurethral manipulation of bulbar urethral strictures is associated with increased stricture complexity and prolonged disease duration. *J Urol.* 187(5):1691-1695
20. Albers P, Fichtner J, Brühl P, Müller SC. Long-term results of internal urethrotomy. *J Urol.* 1996;156:1611-1614.
21. Sachse H. Die internes urethrotomie. *Z Urol Nephrol.* 1980;73: 327-329.

Citation: Yenice MG, Seker KG, Sam E, et al. Comparison of cold-knife optical internal urethrotomy and holmium:YAG laser internal urethrotomy in bulbar urethral strictures. *Cent European J Urol.* 2017; doi: 10.5173/ceju.2017.1391 [Epub ahead of print]

22. Pansodoro V, Emiliozzi P. Internal urethrotomy in the management of anterior urethral strictures: Page 6/11 long term follow-up. *J Urol.* 1996;156:73-75.
23. Naude AM, Heyns CF. What is the place of internal urethrotomy in the treatment of urethral stricture disease?. *Nat Clin Pract Urol.* 2005;2:538-45.
24. Heyns CF, Steenkamp JW, de Kock ML, Whitaker P. Treatment of male urethral strictures: Is repeated dilation or internal urethrotomy useful?. *J Urol.* 1998;160:356-358.
25. Cecen K, Karadag MA, Demir A, Kocaaslan R. PlasmaKinetic™ versus Cold Knife Internal Urethrotomy in Terms of Recurrence Rates: A Prospective Randomized Study. *Urol Int.* 2014;93:460-463.
26. Ozcan L, Polat EC, Otunctemur A, Onen E, Cebeci OO, Memik O, et al. Internal urethrotomy versus plasmakinetic energy for surgical treatment of urethral stricture. *Arch Ital Urol Androl.* 2014; 87:161-164.
27. Tam CA, Elliott SP, Voelzke BB, Myers JB, Vanni AJ, Breyer BN, et al. The International Prostate Symptom Score (IPSS) is an Inadequate Tool to Screen for Urethral Stricture Recurrence After Anterior Urethroplasty. *Urology.* 2016;95:197-201.
28. Tam CA, Voelzke BB, Elliott SP, Myers JB, McClung CD, Vanni AJ, et al. Critical Analysis of the Use of Uroflowmetry for Urethral Stricture Disease Surveillance. *Urology.* 2016;91:197-202.